

V_{ub} , V_{cb} and Decay Constants in Effective Field Theory of Heavy Quarks^a

W.Y. Wang, Y.L. Wu and Y.A. Yan

*Institute of Theoretical Physics, Chinese Academy of Sciences,
Beijing 100080, China*

A new framework of heavy quark effective field theory (HQEFT) is studied and compared with the usual heavy quark effective theory (HQET). $|V_{ub}|$, $|V_{cb}|$ and heavy meson decay constants are extracted in the new framework. HQEFT can yield reasonable results for both exclusive and inclusive decays.

The CKM matrix elements V_{ub} , V_{cb} and the heavy meson decay constants are of crucial importance in particle physics. They have been studied for a long time in the full QCD and in effective theories, but their values are far from being determined precisely. HQET is one of the most favorable effective theories in attacking heavy hadrons. In the usual HQET, one generally decouples the quark and antiquark fields and deals with only one of them independently. This only provides an approximate treatment for some cases though it has been widely applied to various processes.

It has been noticed for a long time that the usual HQET does not always yield consistent and reliable results. The decay constants of heavy mesons receive unexpectedly large $1/m_Q$ order corrections^{1,2}, which makes the calculation based on heavy quark expansion inconsistent, in some sense. For the lifetime ratios of Λ_b baryon and B meson, the theoretical estimate in HQET ($\frac{\tau(\Lambda_b)}{\tau(B^0)} \geq 0.98$ ³) appears to be too large so that it conflicts with the world average of experimental measurements ($\frac{\tau(\Lambda_b)}{\tau(B^0)} = 0.79 \pm 0.05$ ⁴).

In the usual framework of HQET, $1/m_Q$ order corrections to the inclusive decay rate are absent only when the rate is presented in terms of heavy quark mass m_Q rather than the heavy hadron mass m_H ⁵. This seems to be conflict with the case in the exclusive decays where the normalization of the transition matrix elements is given in terms of heavy hadron mass⁶. Such an inconsistency

^aTalk delivered by W.Y. Wang at the International Conference on Flavor Physics (ICFP2001), May 31-June 6, Zhang-Jia-Jie, China.

in the usual HQET may be the main reason that leads to the difficulty in understanding the lifetime difference among the bottom hadrons.

Many processes have not yet been well investigated in HQET. Since the usual HQET does not include the contributions of effective antiquarks in the effective Lagrangian, it is not suitable for analyzing the processes including quark-antiquark pair creation and annihilation. From this point of view, a complete Lagrangian with including both contributions of quarks and antiquarks would be favorable. Transitions such as B decays into excited charmed mesons and heavy-to-light decays have not yet been studied to a satisfactory degree. The large uncertainties require both more precise experimental measurements and a reliable framework in theory.

The heavy quark spin-flavor symmetry can be manifestly exhibited in effective theories. This enables one to analyze different heavy hadrons and processes by using the same set of universal functions, which are independent of the spin-flavor of the heavy quarks. In this way the heavy flavor physics becomes simpler for the reduction of independent parameters. Though the HQET presents a compact framework, there are possibilities of further reducing the number of independent wave functions in the HQEFT as will be illustrated below (see also Refs.^{8,9}).

Strictly speaking, in quantum field theory particle and antiparticle decouple completely only in the limit $m_Q \rightarrow \infty$. To consider the finite quark mass correction precisely, it is necessary to include the contributions from the components of the antiquark fields. One can simply extend the usual HQET to a heavy quark effective field theory with keeping both effective quark and antiquark fields. This was first pointed out by Y. L. Wu in Ref.⁷.

One may start from QCD Lagrangian:

$$\mathcal{L}_{QCD} = \mathcal{L}_{light} + \mathcal{L}_{heavy} = \mathcal{L}_{light} + \bar{Q}(i\not{D} - m_Q)Q. \quad (1)$$

Firstly, denote the quantum field Q as

$$Q = Q^+ + Q^- \quad (2)$$

with Q^+ and Q^- being the two solutions of the Dirac equation, $(i\not{D} - m_Q)Q^\pm = 0$. We decompose the fields $Q^\pm$ into

$$Q^\pm = \hat{Q}_v^\pm + R_v^\pm, \quad (3)$$

where $\hat{Q}_v^\pm \equiv \frac{1 \pm \not{v}}{2} Q^\pm$, $R_v^\pm \equiv \frac{1 \mp \not{v}}{2} Q^\pm$, and introduce new field variables Q_v and $\bar{Q}_v$ defined by

$$Q_v = e^{i\not{p}_Q v \cdot x} \hat{Q}_v, \quad \bar{Q}_v = \bar{\hat{Q}}_v e^{-i\not{p}_Q v \cdot x}. \quad (4)$$

v^μ can be an arbitrary four-vector satisfying $v^2 = 1$. Note that both $\hat{Q}_v$ and R_v contain the components of quark and antiquark fields. $\hat{Q}_v^\pm$ are the ‘large components’ and $R_v^\pm$ the ‘small components’. With above definitions, an effective Lagrangian for the large components ($\hat{Q}_v$ or Q_v) can be easily derived by integrating out the small components $R_v^\pm$:

$$\mathcal{L}_{heavy} \rightarrow \mathcal{L}_{Q,v} = \mathcal{L}_{Q,v}^{++} + \mathcal{L}_{Q,v}^{--} + \mathcal{L}_{Q,v}^{+-} + \mathcal{L}_{Q,v}^{-+} \quad (5)$$

with

$$\mathcal{L}_{Q,v}^{\pm\pm} = \bar{Q}_v^\pm [i\not{D}_\parallel + \frac{1}{2m_Q} i\not{D}_\perp (1 - \frac{i\not{D}_\parallel}{2m_Q})^{-1} i\not{D}_\perp] Q_v^\pm \equiv \bar{Q}_v^\pm A Q_v^\pm, \quad (6)$$

$$\begin{aligned} \mathcal{L}_{Q,v}^{\pm\mp} &= e^{\pm 2im_Q v \cdot x} \bar{Q}_v^\pm [-i\not{D}_\perp + \frac{1}{4m_Q^2} (-i\not{D}_\perp) (1 - \frac{i\not{D}_\parallel}{2m_Q})^{-1} i\not{D}_\perp \\ &\quad \times (1 - \frac{i\not{D}_\parallel}{2m_Q})^{-1} i\not{D}_\perp] Q_v^\mp \equiv e^{\pm 2im_Q v \cdot x} \bar{Q}_v^\pm B Q_v^\mp. \end{aligned} \quad (7)$$

In the usual HQET, quark and antiquark fields are completely decoupled, and the effective Lagrangian there includes only the Q^+ component in Eq.(2). As a result, the effective Lagrangian in HQET contains only $\mathcal{L}_{Q,v}^{++}$ but not the other 3 parts in Eq.(5).

If further integrating out the antiquark fields (but not neglecting their contributions), the effective Lagrangian can be written in the following form in terms of only quark fields:

$$\begin{aligned} \mathcal{L}_{eff} &= \bar{Q}_v^+ \{ i\not{D}_\parallel + \frac{1}{2m_Q} i\not{D}_\perp (1 - \frac{i\not{D}_\parallel}{2m_Q})^{-1} i\not{D}_\perp - [i\not{D}_\perp + \frac{1}{2m_Q} i\not{D}_\perp \\ &\quad \times (1 - \frac{i\not{D}_\parallel}{2m_Q})^{-1} i\not{D}_\perp \frac{1}{i\not{D}_\parallel} i\not{D}_\perp] [i\not{D}_\parallel - 2m_Q - i\not{D}_\perp \frac{1}{i\not{D}_\parallel} i\not{D}_\perp]^{-1} \\ &\quad \times [i\not{D}_\perp + \frac{1}{2m_Q} i\not{D}_\perp \frac{1}{i\not{D}_\parallel} i\not{D}_\perp (1 - \frac{i\not{D}_\parallel}{2m_Q})^{-1} i\not{D}_\perp] \} Q_v^+. \end{aligned} \quad (8)$$

When the mass of a heavy quark is much larger than the QCD scale Λ_{QCD} , this effective Lagrangian can be expanded in inverse power of the heavy quark mass and be straightforwardly written as $\mathcal{L}_{eff} = \mathcal{L}_{eff}^{(0)} + \mathcal{L}_{eff}^{(1/m_Q)}$ with

$$\mathcal{L}_{eff}^{(0)} = \bar{Q}_v^+ (i\not{D}_\parallel) Q_v^+, \quad (9)$$

$$\begin{aligned} \mathcal{L}_{eff}^{(1/m_Q)} &= \frac{1}{m_Q} \bar{Q}_v^+ (i\not{D}_\perp)^2 Q_v^+ + \frac{1}{2m_Q^2} \bar{Q}_v^+ i\not{D}_\perp (i\not{D}_\parallel) i\not{D}_\perp Q_v^+ \\ &\quad + \frac{1}{4m_Q^2} \bar{Q}_v^+ (i\not{D}_\perp)^2 \frac{1}{i\not{D}_\parallel} (i\not{D}_\perp)^2 Q_v^+ + O(\frac{1}{m_Q^3}), \end{aligned} \quad (10)$$

To show the contributions of antiquarks more clearly, the Lagrangian can be written as:

$$\mathcal{L} \rightarrow \mathcal{L}_{Q,v} \rightarrow \mathcal{L}_{eff} = \mathcal{L}_{Q,v}^{++} + \tilde{\mathcal{L}}_{Q,v}^{++}, \quad (11)$$

$$\begin{aligned} \tilde{\mathcal{L}}_{Q,v}^{++} &\equiv \mathcal{L}_{Q,v}^{+-} + \mathcal{L}_{Q,v}^{-+} + \mathcal{L}_{Q,v}^{--} \\ &= \frac{1}{2m_Q} \bar{Q}_v^+ A \frac{1}{i\not{D}_{\parallel}} i\not{D}_{\perp} [1 - \frac{i\not{D}_{\parallel}}{2m_Q} + \frac{1}{2m_Q} i\not{D}_{\perp} \frac{1}{i\not{D}_{\parallel}} i\not{D}_{\perp}]^{-1} i\not{D}_{\perp} \frac{1}{i\not{D}_{\parallel}} A Q_v^+. \end{aligned} \quad (12)$$

$\tilde{\mathcal{L}}_{Q,v}^{++}$ is the additional contributions to the part $\mathcal{L}_{Q,v}^{++}$ that is adopted in the usual HQET. This additional part may be regarded as the effective potential of heavy quark due to the exchanges of virtual antiquarks. When one imposes the on-shell condition $AQ_v^+ = 0$, i.e. $\mathcal{L}_{Q,v}^{++} = 0$, the effective potential $\tilde{\mathcal{L}}_{Q,v}^{++}$ also vanishes. For an off-shell heavy quark in a hadron, $\tilde{\mathcal{L}}_{Q,v}^{++} \neq 0$, and the leading order contribution of the effective potential is $\tilde{\mathcal{L}}_{Q,v}^{++}|_{LO} = \bar{Q}_v^+ \frac{(i\not{D}_{\perp})^2}{2m_Q} Q_v^+$. As is expected, antiquarks contribute from the order of $1/m_Q$.

The effective Lagrangian $\mathcal{L}_{eff}$ automatically preserves the velocity reparametrization invariance as well as Lorentz invariance without the need of summing over the velocity. This can be understood easily by noticing the fact $\mathcal{L}_{QCD} = \mathcal{L}_{light} + \mathcal{L}_{Q,v} = \mathcal{L}_{light} + \mathcal{L}_{Q,v'}$.

It is also noticed that Luke's theorem is a natural result of this HQEFT, without the need of on-shell condition $iv \cdot DQ_v^+ = 0$ ⁸. This result is nontrivial and has its importance in explaining the bottom hadron lifetime ratios in HQEFT.

Furthermore, fewer independent functions (parameters) are needed in HQEFT than in HQET. For example, to characterize the $1/m_Q$ order corrections to heavy-to-heavy transition matrix elements, HQET needs all together 8 functions, while HQEFT needs only 3. And for the $1/m_Q$ order corrections to heavy meson decay constants, HQET needs 4 parameters, while HQEFT needs only 2. This simplification has been illustrated in detail in Refs.^{8,9}.

In the HQEFT, heavy meson decay constants can be written as⁹

$$f_M = \frac{F}{\sqrt{m_M}} \{1 + \frac{1}{m_Q} (g_1 + 2d_M g_2)\}. \quad (13)$$

F is the leading order parameter, while g_1, g_2 characterize the $1/m_Q$ order corrections. These have been calculated using sum rule methods with the results⁹: $\bar{\Lambda} = 0.53 \pm 0.08 \text{ GeV}$, $F = 0.30 \pm 0.06 \text{ GeV}^{3/2}$, $g_1 = 0.54 \pm 0.12 \text{ GeV}$, $g_2 = -0.06 \pm 0.02 \text{ GeV}$. Notice that $|g_1|/m_Q$ remains small enough so that the $1/m_Q$ expansion in the new framework of HQEFT appears to be reliable. The scaling law $f_M \sim F/\sqrt{m_M}$ is only slightly broken at the $1/m_Q$ order.

As a comparison, decay constants in the usual HQET is represented as

$$f_M = \frac{F}{\sqrt{m_M}} \left(1 - d_M \frac{\bar{\Lambda}}{6m_Q} + \frac{G_1}{m_Q} + d_M \frac{G_2}{m_Q} \right). \quad (14)$$

with the $1/m_Q$ order corrections very large ($G_1 \approx -2.0\text{GeV}$ in Ref.¹, $G_1 \approx -0.8\text{GeV}$ in Ref.²). So, in the HQET, the scaling law is seriously broken.

For the decay constants, we get in HQEFT when no QCD radiative corrections are considered⁹: $f_B(m_b) = 159 \pm 42$ MeV, $f_{B^*}(m_b) = 166 \pm 38$ MeV, $f_D(m_c) = 251 \pm 99$ MeV, $f_{D^*}(m_c) = 293 \pm 86$ MeV. With two-loop QCD corrections, we get: $f_B(m_b) = 196 \pm 44$ MeV, $f_{B^*}(m_b) = 206 \pm 39$ MeV, $f_D(m_c) = 298 \pm 109$ MeV, $f_{D^*}(m_c) = 354 \pm 90$ MeV. These agree with results obtained from lattice simulations or other phenomenological approaches¹⁰.

Semileptonic decays $B \rightarrow D^*(D)l\nu$ provide one of the main approaches to extract the value of $|V_{cb}|$. The differential decay rates are:

$$\begin{aligned} \frac{d\Gamma(B \rightarrow D^*l\nu)}{d\omega} &= \frac{G_F^2}{48\pi^3} (m_B - m_{D^*})^2 m_{D^*}^3 \sqrt{\omega^2 - 1} (\omega + 1)^2 \\ &\times \left[1 + \frac{4\omega}{\omega + 1} \frac{m_B^2 - 2\omega m_B m_{D^*} + m_{D^*}^2}{(m_B - m_{D^*})^2} \right] |V_{cb}|^2 \mathcal{F}^2(\omega), \end{aligned} \quad (15)$$

$$\frac{d\Gamma(B \rightarrow Dl\nu)}{d\omega} = \frac{G_F^2}{48\pi^3} (m_B + m_D)^2 m_D^3 (\omega^2 - 1)^{3/2} |V_{cb}|^2 \mathcal{G}^2(\omega) \quad (16)$$

with $\omega = v \cdot v'$, $\mathcal{F}(\omega) = \eta_A h_{A_1}(\omega)$, $\mathcal{G}(\omega) = \eta_V [h_+(\omega) - \frac{m_B - m_D}{m_B + m_D} h_-(\omega)]$, where $|V_{cb}| \mathcal{F}(1) = 0.0352 \pm 0.0026$, $|V_{cb}| \mathcal{G}(1) = 0.0386 \pm 0.0041$ ¹¹; $\eta_A = 0.960 \pm 0.007$, $\eta_V = 1.022 \pm 0.004$ ¹².

Due to the Luke's theorem, the factors h_{A_1} , h_+ and h_- are protected from $1/m_Q$ order corrections at the zero recoil point $\omega = 1$. For this reason, in the usual HQET, the $B \rightarrow D^*l\nu$ decay is more favorable for the extraction of $|V_{cb}|$, because its decay rate receives no corrections of $1/m_Q$ order. In the new framework of HQEFT, however, $h_-(\omega) = 0$ ⁸. This implies that both the differential decay rates of channels $B \rightarrow Dl\nu$ and $B \rightarrow D^*l\nu$ receive no order $1/m_Q$ corrections. Therefore both channels can be used to extract $|V_{cb}|$ reliably.

The HQEFT gives interesting relations between meson masses and wave functions⁸. Those relations enable one to get the zero recoil values of HQEFT wave functions, and furthermore to extract $|V_{cb}|$ from the meson masses. We get $|V_{cb}| = (3.78 \pm 0.28_{\text{exp}} \pm 0.18_{\text{th}}) \times 10^{-2}$ from $B \rightarrow D^*l\nu$ decay, and $|V_{cb}| = (3.82 \pm 0.41_{\text{exp}} \pm 0.28_{\text{th}}) \times 10^{-2}$ from $B \rightarrow Dl\nu$ decay⁸, which are close to each other, and agree with the world average¹³: $(3.67 \pm 0.23_{\text{exp}} \pm 0.18_{\text{th}}) \times 10^{-2}$ with using dispersion relation $\mathcal{F}(\omega)$ parameterization, and $(3.92 \pm 0.30_{\text{exp}} \pm 0.19_{\text{th}}) \times 10^{-2}$ with using linear $\mathcal{F}(\omega)$ parameterization.

Semileptonic decays $B \rightarrow \pi(\rho)l\nu$ provide two of the most direct channels to estimate $|V_{ub}|$. One can parameterize the leading order transition matrix elements in the effective theory of heavy quarks as^{14,15}

$$\begin{aligned}\langle \pi(p) | \bar{u} \Gamma Q_v^+ | B_v \rangle &= -Tr[\pi(v, p) \Gamma \mathcal{M}(v)], \\ \langle \rho(p, \epsilon^*) | \bar{u} \Gamma Q_v^+ | B_v \rangle &= -iTr[\Omega(v, p) \Gamma \mathcal{M}_v],\end{aligned}$$

where

$$\begin{aligned}\pi(v, p) &= \gamma^5[A(v \cdot p, \mu) + \not{p}B(v \cdot p, \mu)], \\ \Omega(v, p) &= L_1(v \cdot p)\not{\epsilon}^* + L_2(v \cdot p)(v \cdot \epsilon^*) + [L_3(v \cdot p)\not{\epsilon}^* + L_4(v \cdot p)(v \cdot \epsilon^*)]\not{p}\end{aligned}\quad (17)$$

with $\hat{p}^\mu = \frac{p^\mu}{v \cdot p}$. A , B and L_i ($i = 1, 2, 3, 4$) are the leading order wave functions characterizing the heavy-to-light matrix elements in the effective theory. In heavy-to-light decays, the HQS loses some of its predictive power. However, HQS and relevant effective theories are still useful since they give us relations between different channels. For example, $B \rightarrow \rho l \nu$ and $D \rightarrow \rho l \nu$ are characterized by the same set of wave functions L_i . We calculated these leading order wave functions using light cone sum rules. Consider the π distribution functions up to twist 4 and the ρ distribution functions up to twist 2, we get $|V_{ub}| = (3.4 \pm 0.5_{\text{exp}} \pm 0.5_{\text{th}}) \times 10^{-3}$ from $B \rightarrow \pi l \nu$ decay¹⁶, and $|V_{ub}| = (3.9 \pm 0.6_{\text{exp}} \pm 0.7_{\text{th}}) \times 10^{-3}$ from $B \rightarrow \rho l \nu$ decay¹⁷. These leading order results agree well with those from full QCD calculations¹⁸: $(3.9 \pm 0.6_{\text{exp}} \pm 0.6_{\text{th}}) \times 10^{-3}$ from $B \rightarrow \pi l \nu$ decay, or $(3.4 \pm 0.6_{\text{exp}} \pm 0.5_{\text{th}}) \times 10^{-3}$ from $B \rightarrow \rho l \nu$ decay, and are also compatible with the combined result from the analyses based on different models and treatments on $B \rightarrow \pi(\rho)l\nu$ transitions¹⁹: $|V_{ub}| = (3.25 \pm 0.14_{-0.29}^{+0.21} \pm 0.55) \times 10^{-3}$.

Let us also take a look on the application of HQEFT on inclusive decays. The HQET predictions on the lifetime differences of bottom hadrons are³: $\frac{\tau(B^0)}{\tau(B^{\pm})} \approx 1$, $\frac{\tau(\Lambda_b)}{\tau(B^0)} \geq 0.98$, while the world average values of experiments are⁴: $\frac{\tau(B^0)}{\tau(B^{\pm})} = 0.94 \pm 0.04$, $\frac{\tau(\Lambda_b)}{\tau(B^0)} = 0.79 \pm 0.05$. This is a well-known contradiction in heavy flavor physics. Calculations^{20,21} in HQEFT give $\frac{\tau(B_s^0)}{\tau(B^0)} = 0.96 \pm 0.06$, $\frac{\tau(\Lambda_b)}{\tau(B^0)} = 0.78 \pm 0.05$, in good agreement with the experimental values.

Calculations in HQEFT on inclusive semileptonic decay rates also lead to $|V_{cb}|$ and $|V_{ub}|$ values^{20,21}: $|V_{cb}| = (3.89 \pm 0.05_{\text{exp}} \pm 0.20_{\text{th}}) \times 10^{-2}$, $|V_{ub}| = (3.48 \pm 0.62_{\text{exp}} \pm 0.11_{\text{th}}) \times 10^{-3}$, which agree well with those obtained from exclusive decays.

For the charm counting in B^0 decays, HQEFT predicts^{20,21} $n_c = 1.19 \pm 0.04$, which is smaller than predictions in HQET²²: $n_c = 1.24 \pm 0.06$ in OS scheme, and $n_c = 1.30 \pm 0.07$ in $\overline{MS}$ scheme. So the value of n_c obtained in

HQEFT is, compared to the HQET prediction, closer to the world average²³: $n_c = 1.17 \pm 0.04$.

In these applications of HQEFT into inclusive decays, the mass entering into the inclusive decay rates is the ‘dressed heavy quark’ mass $\hat{m}_Q^5$ instead of the quark mass m_Q^5 , where

$$\hat{m}_Q = m_Q + \bar{\Lambda} = m_{H_Q}(1 + O(1/m_{H_Q}^2)) \quad (18)$$

with $\bar{\Lambda}$ being the binding energy of the heavy hadron. And therefore hadrons have different phase space effects in HQEFT. These are the main reasons for the improvement of the new HQEFT with respect to the usual HQET in studying inclusive decays.

As a summary, the HQEFT gives reasonable results for $|V_{ub}|$, $|V_{cb}|$ and heavy meson decay constants. In all discussed applications in exclusive and inclusive decays, the HQEFT proves to be consistent, reliable and simple. We found that the antiquark components do contribute measurable effects, which in turn supports the inclusion of both quark and antiquark contributions in the effective Lagrangian. The HQEFT should be able to be applied to many other processes of heavy hadrons, and further improve our understanding on the heavy flavor physics.

Acknowledgement This work was supported in part by Chinese Academy of Sciences and by NSFC under grant # 19625514.

1. M. Neubert, Phys. Rev. D **46**, 1076 (1992).
2. P. Ball and V. M. Braun, Nucl. Phys. B **421**, 593 (1994).
3. H. Y. Cheng and K. C. Yang, Phys. Rev. D **59**, 014011 (1999).
4. For updated world averages of B hadron lifetimes and lifetime difference, see J. Alcaraz et. al. (LEP B Lifetime Group), <http://wwwcn.cern.ch/claibes/lepblife.html>.
5. J. Chay, H. Georgi, and B. Grinstein, Phys. Lett. **B247** 399 (1990).
6. M. E. Luke, Phys. Lett. B **252**, 447 (1990).
7. Y. L. Wu, Modern Phys. Lett. A **8**, 819 (1993).
8. W. Y. Wang, Y. L. Wu and Y. A. Yan, Int. J. Mod. Phys. A **15**, 1817 (2000).
9. W. Y. Wang and Y. L. Wu, Int. J. Mod. Phys. A **16**, 377 (2001).
10. D. Becirevic, et. al., BUHEP-00-3, hep-lat/0002025; L. Lellouch and C. D. Lin, CERN-TH/99-344, hep-ph/9912322; C. Bernard et. al., Nucl. Phys. Proc. Suppl. **83**, 289 (2000); S. Collins, et. al., Phys. Rev. D **60**, 074504 (1999).
11. M. Neubert, CERN-TH/98-2, hep-ph/9801269.
12. A. Czarnecki, Phys. Rev. Lett. **22**, 4124 (1996).
13. Particle Data Group, Eur. Phys. J. C **15**, 2000.

14. G. Burdman, Z. Ligeti, M. Neubert and Y. Nir, Phys. Rev. D **49**, 2331 (1994).
15. C. S. Huang, C. Liu and C. T. Yan, Phys.Rev. D **62**, 054019 (2000).
16. W. Y. Wang and Y. L. Wu, Phys. Lett. B **515**, 57 (2001).
17. W. Y. Wang and Y. L. Wu, Phys. Lett. B **519**, 219 (2001).
18. A. Khodjamirian and R. Rückl, WUE-ITP-97-049, MPI-PhT/97-85, hep-ph/9801443.
19. B. H. Behrens et. al.(CLEO Collab.), Phys. Rev. D **61**, 052001 (2000).
20. Y. A. Yan, Y. L. Wu and W. Y. Wang, Int. J. Mod. Phys. A **15**, 2735 (2000).
21. Y. L. Wu and Y. A. Yan, Int. J. Mod. Phys. A **16**, 285 (2001).
22. E. Bagan, P. Ball, M. Braun and P. Gosdzinsky, Nucl. Phys. B **432**, 3 (1994); Phys. Lett. B **342**, 362 (1995); Erratum, ibid. B **374**, 363 (1996); E. Bagan, P. Ball, B. Fiol and P. Gosdzinsky, Phys. Lett. B **351**, 546 (1995).
23. Particle Data Group, Eur. Phys. J. C **3**, 1998.